

Perinatal Mood Disorder Research Articles

Abramowitz JS, et al. (2003). Obsessive-compulsive symptoms in pregnancy and the puerperium: A review of the literature. *Anxiety Disorders*, 17: 461-478.

Affonso, D.D., De, A.K., Andrews Horowitz, J.A., et al. (2000). An international study exploring levels of postpartum depressive symptomatology. *Journal of Psychosomatic Research*, 49, 207-216.

Allister, L., et al. (2001). The Effects of Maternal Depression on Fetal Heart Rate Response to Vibroacoustic Stimulation. *Developmental Neuropsychology*, 20: 639-651.

Appleby, L., Warner, R., Whitton, A., et al. (1997). A controlled study of fluoxetine and cognitive-behavioural counselling in the treatment of postnatal depression. *British Medical Journal*, 314, 932-936.

Appleby, L., Koren, G., & Sharp, D. (1999). Depression in pregnant and postnatal women: an evidence-based approach to treatment in primary care. *British Journal of General Practice*, 780-782.

Armstrong, K.L., Fraser, J.A., Dadds, M.R., & Morris, J. (1999). A randomized, controlled trial of nurse home visiting to vulnerable families with newborns. *Journal of Pediatric Child Health*, 35: 237-244.

Austin, M.P., et al. (2005). Prenatal stress, the hypothalamic–pituitary–adrenal axis, and fetal and infant neurobehaviour. *Early Human Development*, 81: 917-926.

Barnet, B., Duggan, A.K., Wilson, M.D., & Joffe, A. (1995). Association between postpartum substance use and depressive symptoms, stress, and social support in adolescent mothers. *Pediatrics*, 96: 659-666.

Beck, C.T. (2004). Birth Trauma: in the Eye of the Beholder, *Nursing Research*, 53: 28-35.

Beck, C.T. (2003). Recognizing and Screening for Postpartum in Moms of NICU infants. *Advanced Neonatal Care*. 3: 1.

Bennet H et al, (2004). Depression during Pregnancy, Overview of Clinical Factors. *Clinical Drug Investigations*. 24: 157-17.

Bledsoe, S.E., Grote, N.K. (2006). Treating Depression During Pregnancy and the Postpartum: A Preliminary Meta-Analysis. *Research on Social Work Practice*. 16; 109–120.

Berggren-Clive, K. (1998). Out of the darkness and into the light: Women's experiences with depression after childbirth. *Canadian Journal of Community Mental Health*, 17: 103-120.

Bewley, C. (1999). Postnatal depression. *Nursing Standard*, 13: 49-56.

Borrill, J. (1998). Detecting and preventing postnatal depression. *Community Nurse*, 19-20.

Brennan, P.A., Andersen, M.J, Hammen, C., et. Al. (2000). Chronicity, severity, and timing of maternal depressive symptoms: relationships with child outcomes at age 5. *Developmental Psychology*, 36: 759-766.

Bagedahl-Strindlund, M. & Monsen, B.K. (1998). Postnatal depression: A hidden illness. *Acta Psychiatria Scandinavia*, 98, 272-275.

Choi Y, Bishai D, & Minkovitz, C, (2009). Multiple Births are a Risk Factor for Postpartum Maternal Depressive Symptoms. *Pediatrics*, 123:1147-1154.

Coral, M., Kuan, A., & Kostaras, D. (2000). Bright light therapy effect on postpartum depression. *American Journal of Psychiatry*, 157: 303-304.

Corbin, A. & O'Grady, J. (2002). Tackling postnatal depression. *Professional Nurse*, 17, 372-373.

Cox, J.L., Holden, J.M., & Sagovsky, R. (1987). Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry*, 150, 782-786.

Dennis, CL & McQueen K. (2009). The Relationship between infant-feeding outcomes and postpartum depression. *Pediatrics*, 123: 736-751.

De Mier, R.L. & Hynan M.T. (2000). A measurement model of perinatal stressors: Identifying risk for postnatal emotional distress in mothers of high-risk infants. *Journal of Clinical Psychology*, 56: 89-100.

Diego, M. et al (2002). Facial expression & EEG in infants of intrusive and withdrawn mothers with depressive symptoms. *Depression & Anxiety*, 15: 107

Edebohls, L. & Ecklund, C. (2002). Commentary on Andrea Yates, Postpartum depression: Practical advice from two nurse practitioners. *Pediatric Nursing*, 22: 298-299.

Elliot, S.A., Leverton, T.J., Sanjack, M., et al. (2000). Promoting mental health after childbirth: A controlled trial of primary prevention of postnatal depression. *British Journal of Clinical Psychology*, 39, 223-241.

Epperson, N., Czarkowski, K.A., Ward-O'Brien, D., et. al. (2001). Maternal sertraline treatment and serotonin transport in breast-feeding mother-infant pairs. *American Journal of Psychiatry*, 158: 1631-1637.

Epperson CN, Jatlow PI, Czarkowski K, et al. (2003). Maternal fluoxetine treatment in the postpartum period: effects on platelet serotonin and plasma drug levels in breastfeeding mother-infant pairs. *Pediatrics*, 112:425.

Field, T., Grizzle, N., Scafidi, F., & Schanberg, S. (1996). Massage and relaxation therapies, effects on depressed adolescent mothers. *Adolescence*, 34: 904-911.

Field T. Early interventions for infants of depressed mothers, *Pediatrics*. 2005.102:1305

Hale, T. (2002). *Medications and Mothers' Milk*. 10th Edition. Pharmasoft Publishing.

Hall, L.A., Kotch, J.B., Brown, D., et al. (1996). Self-esteem as a mediator of the effects of stressors and social on depressive symptoms in postpartum mothers. *Nursing Research*, 45: 231-238.

Hayes, B.A. Muller, R., & Bradley B.S. (2001). Perinatal depression: A randomized controlled trial of an antenatal education intervention for primiparas. *Birth*, 28, 28-35.

Hayes, M.J., Roberts, S., & Davare, A. (2000). Transactional conflict between psychobiology and culture in the etiology of postpartum depression. *Medical Hypotheses*, 55: 266-276.

Johnston, S.J., Boyce, P.M., Hickey, A.R., et al. (2001). Obstetric risk factors for postnatal depression in urban and rural community samples. *Australian & New Zealand Journal of Psychiatry*, 35, 69-74.

Jones, H.W. & Venis, J.A. (2001). Identification and classification of postpartum psychiatric disorders. *Journal of Psychosocial Nursing*, 39, 23-47.

Josefsson, A., Angelsioo, L., Berg, G., et al. (2002). Obstetric, somatic, and demographic risk factors for postpartum depressive symptoms. *Obstetrics and Gynecology*, 99, 223-228.

Kendall-Tackett. (2001). A new paradigm for depression in new mothers, *Intl Breastfeeding Journal*, 2:6.

Kleiman, K.R. & Raskin, V.D. (1994) *This Isn't What I Expected: Overcoming Postpartum Depression*, New York: Bantam Books.

Lane, B., Roufeil, L.M. Williams, S., et al. (2002). It's just different in the country: Postnatal depression and group therapy in a rural setting. *Social Work Health and Mental Health: Practices, Research and Programs*, 333-348.

Lawrie, T.A., Herxheimer, A., & Dalton, K. (2002). Oestrogens and progestogens for preventing and treating postnatal depression. *The Cochrane Library*, 4. Available: www.cochranelibrary.com.

Levinson-Castiel R, Merlob P, Linder N, et al. (2006). Neonatal abstinence syndrome after in utero exposure to selective serotonin reuptake inhibitors in term infants. *Archives Pediatric Adolescent Medicine*, 160:173–176.

Lindahl V, Pearson JL, & Colpe L. (2005). Prevalence of suicidality during pregnancy and the postpartum. *Archives Women's Mental Health*. 8, 77-87.

Lindgren, K. (2001). Relationships among maternal-fetal attachment, prenatal depression, and health practices in pregnancy. *Research in Nursing & Health*, 24, 203-217.

Nermeroff, C. (2001). Progress in the battle with the black dog: Advances in the treatment of depression. *American Journal Psychiatry*, 158: 1555-1557.

Misri, S., Kostaras, X., Fox, D., et al. (2000). The impact of partner support in the treatment of postpartum depression. *Canadian Journal Psychiatry*. 45: 554-558.

Misri S, Sinclair CA, Kuan AJ, (1997). Breast-feeding and postpartum depression: Is there a relationship? *Canadian Journal Psychiatry*, 42:1061-5.

Newport, J.D, Hosteter, A., Arnold, A., et al. (2002). The treatment of postpartum depression: Minimizing infant exposures. *Journal of Clinical Psychiatry*, 63, 31-44.

Oberlander et. al. (2008). Infant serotonin transporter (SLC6A4) promoter genotype is associated with adverse neonatal outcomes after prenatal exposure to serotonin reuptake inhibitor medications. *Molecular Psychiatry*, 13, 65–73.

O'Hara, M.W., et al. (2000). Efficacy of Interpersonal Psychotherapy for Postpartum Depression. *Archives of General Psychiatry*, 59: 1039-1054.

Paulson, Daubner and Leiferman. (2006). Individual and combined effects of postpartum depression in mothers and fathers on parenting behavior. *Pediatrics*, 118: 659-668.

Paulson, et al. (2006). Individual and Combined Effects of Postpartum Depression in Mothers and Fathers on Parenting Behavior. *Pediatrics*, 118: 659-668.

Seguin, L., Potvin, L., St-Denis, M., et al (1999). Depressive symptoms in late postpartum among low socioeconomic status women. *Birth*, 26: 157-163.

Shepard. C. & Feifler, B.V. (1995). Postpartum psychiatric disorders: A guide for the practicing physician. *North Carolina Medical Journal*, 56: 386-388.

Strass, P. (2002). Rural community health: Postpartum depression support. *Canadian Nurse*. 98: 25-28.

Steiner, M. (1998). Perinatal mood disorders: Position paper. *Psychopharmacology Bulletin*, 34: 301-306.

Tatano Beck, C. (2001). Predictors of postpartum depression: An update. *Nursing Research*, 50: 275-285.

Tarkka, M-T. & Paunonen, M. (1996). Social support provided by nurses to recent mothers on a maternity ward. *Journal of Advanced Nursing*, 23, 1202-1206.

Warner, R., Appleby, L., Whitton, A., et al. (1996). Demographic and obstetric risk factors for postnatal psychiatric morbidity. *British Journal of Psychiatry*, 168, 607-611.

Winans, E.A. (2001). Antidepressant use during lactation. *Journal of Human Lactation*, 17: 256-261.

Wisner, K.L. (1999). Prevention and treatment of postpartum mood disorders. *The Cochrane Library*, 4. Available: www.cochranelibrary.com.

Wisner et al. (2004). Prevention of Postpartum Depression: A Randomized Pilot. *American Journal of Psychiatry*, 161: 1290-1292.

Zambaldi CF, Cantilino A, Montenegro AC, et al. (2009). Postpartum obsessive-compulsive disorder: prevalence and clinical characteristics. *Comprehensive Psychiatry*, 50:503-509.

Zlotnick, C., Johnson, S.L., Miller, I.W., et al. (2001). Postpartum depression in women receiving public assistance: Pilot study of an interpersonal therapy oriented group intervention. *American Journal of Psychiatry*, 158: 638-640.